

# ОБ ОДНОМ КЛАССЕ ПОДРЕШЕТОК РЕШЕТКИ ПОДГРУПП КОНЕЧНОЙ ГРУППЫ

Н.С. Косенок, И.В. Блинец

*Гомельский государственный университет имени Франциска Скорины*

# ON A CLASS OF SUBLATTICES OF THE SUBGROUP LATTICE OF A FINITE GROUP

N.S. Kosenok, I.V. Blisnets

*Francisk Skorina Gomel State University*

**Аннотация.** В данной работе:  $G$  – конечная группа;  $\sigma = \{\sigma_i \mid i \in I\}$  – некоторое разбиение множества всех простых чисел  $\mathbb{P}$ ;  $\Pi \subseteq \sigma$ ;  $\sigma(n) = \{\sigma_i \mid \sigma_i \cap \pi(n) \neq \emptyset\}$  ( $n$  – целое число) и  $\sigma(G) = \sigma(|G|)$ . Группа  $G$  называется: (i)  $\sigma$ -примарной, если  $G$  является  $\sigma_i$ -группой для некоторого  $i \in I$ ; (ii)  $\sigma$ -нильпотентной, если  $G$  – прямое произведение  $\sigma$ -примарных групп;  $\Pi$ -группой, если  $\sigma(G) \subseteq \Pi$ . Подгруппа  $A$  конечной группы  $G$  называется: (i)  $\sigma$ -субнормальной в  $G$ , если в  $G$  существует цепь подгрупп  $A = A_0 \leq A_1 \leq \dots \leq A_t = G$  такая, что либо  $A_{i-1} \trianglelefteq A_i$ , либо  $A_i / (A_{i-1})_{A_i}$  является  $\sigma$ -примарной группой для всех  $i = 1, \dots, t$ ; (ii) холловской  $\Pi$ -подгруппой  $G$ , если  $A$  является  $\Pi$ -группой и  $\sigma(|G:A|) \cap \Pi = \emptyset$ . Мы говорим, что подгруппа  $H$  группы  $G$  является строго  $\sigma$ -субнормальной, если  $H^G / H_G$  является  $\sigma$ -нильпотентной группой. В данной работе мы доказываем, что множество всех строго  $\sigma$ -субнормальных подгрупп, перестановочных с холловой  $\Pi$ -подгруппой конечной группы  $G$ , образует подрешётку решётки всех подгрупп  $L(G)$  группы  $G$ .

**Ключевые слова:** конечная группа, решетка подгрупп, группа операторов, подрешетка решетки, холлова  $\Pi$ -подгруппа.

**Для цитирования:** Косенок, Н.С. Об одном классе подрешеток решетки подгрупп конечной группы / Н.С. Косенок, И.В. Блинец // Проблемы физики, математики и техники. – 2025. – № 4 (65). – С. 72–74. – DOI: [https://doi.org/10.54341/20778708\\_2025\\_4\\_65\\_72](https://doi.org/10.54341/20778708_2025_4_65_72). – EDN: KEQNCF

**Abstract.** In this paper:  $G$  is a finite group;  $\sigma = \{\sigma_i \mid i \in I\}$  is some partition of the set of all primes  $\mathbb{P}$ ;  $\Pi \subseteq \sigma$ ;  $\sigma(n) = \{\sigma_i \mid \sigma_i \cap \pi(n) \neq \emptyset\}$  ( $n$  is an integer) and  $\sigma(G) = \sigma(|G|)$ . A group  $G$  is said to be: (i)  $\sigma$ -primary provided  $G$  is a  $\sigma_i$ -group for some  $i \in I$ ; (ii)  $\sigma$ -nilpotent if  $G$  is the direct product of  $\sigma$ -primary groups; a  $\Pi$ -group if  $\sigma(G) \subseteq \Pi$ . A subgroup  $A$  of a finite group  $G$  is said to be: (i)  $\sigma$ -subnormal in  $G$  if there is a subgroup chain  $A = A_0 \leq A_1 \leq \dots \leq A_t = G$  such that either  $A_{i-1} \trianglelefteq A_i$  or  $A_i / (A_{i-1})_{A_i}$  is  $\sigma$ -primary for all  $i = 1, \dots, t$ ; (ii) a Hall  $\Pi$ -subgroup of  $G$  if  $A$  is a  $\Pi$ -group and  $\sigma(|G:A|) \cap \Pi = \emptyset$ .

We say that a subgroup  $H$  of  $G$  is strongly  $\sigma$ -subnormal if  $H^G / H_G$  is  $\sigma$ -nilpotent. In this paper, we prove that the set of all strongly  $\sigma$ -subnormal subgroups which permute with a Hall  $\Pi$ -subgroup of a finite group  $G$  forms a sublattice of the lattice of all subgroups  $L(G)$  of  $G$ .

**Keywords:** finite group, lattice of subgroups, operator group, sublattice of a lattice, Hall  $\Pi$ -subgroup.

**For citation:** Kosenok, N.S. On a class of sublattices of the subgroup lattice of a finite group / N.S. Kosenok, I.V. Blisnets // Problems of Physics, Mathematics and Technics. – 2025. – № 4 (65). – P. 72–74. – DOI: [https://doi.org/10.54341/20778708\\_2025\\_4\\_65\\_72](https://doi.org/10.54341/20778708_2025_4_65_72). – EDN: KEQNCF

## Introduction

Throughout this paper, all groups are finite and  $G$  always denotes a finite group. Moreover,  $\mathbb{P}$  is the set of all primes and  $\sigma = \{\sigma_i \mid i \in I\}$  is some partition of  $\mathbb{P}$ ;  $\Pi \subseteq \sigma$  and  $\Pi' = \sigma \setminus \Pi$ .

If  $n$  is an integer, the symbol  $\pi(n)$  denotes the set of all primes dividing  $n$ ; as usual,  $\pi(G) = \pi(|G|)$ , the set of all primes dividing the

order of  $G$ ;  $\sigma(n) = \{\sigma_i \mid \sigma_i \cap \pi(n) \neq \emptyset\}$  and  $\sigma(G) = \sigma(|G|)$  [1]–[4].

A  $\sigma$ -property of a group [1]–[4] is understood to be any of its properties that depends on  $\sigma$  and which does not imply any restrictions on  $\sigma$ .

Before continuing, let us recall some of the most important concepts of the theory of  $\sigma$ -properties of a group.

A group  $G$  is said to be [1]–[4]:

- (i)  $\sigma$ -primary if  $G$  is a  $\sigma_i$ -group for some  $i \in I$ ;
- (ii)  $\sigma$ -nilpotent if  $G$  is the direct product of  $\sigma$ -primary groups;

A subgroup  $A$  of  $G$  is said to be [1]–[4]:

- (i)  $\sigma$ -subnormal in  $G$  if there is a subgroup chain

$$A = A_0 \leq A_1 \leq \dots \leq A_n = G$$

such that either  $A_{i-1} \trianglelefteq A_i$  or  $A_i / (A_{i-1})_{A_i}$  is  $\sigma$ -primary for all  $i = 1, \dots, n$ ;

- (ii)  $\sigma$ -permutable in  $G$  provided  $G$  is  $\sigma$ -full, that is,  $G$  has a Hall  $\sigma_i$ -subgroup for all  $i \in I$  and  $A$  permutes with all Hall  $\sigma_i$ -subgroups of  $G$  for all  $i$ .

We use  $H^G$  to denote the normal closure of the subgroup  $H$  in  $G$  (so  $H^G$  is the intersection of all normal subgroups of  $G$  containing  $H$ ), and  $H_G$  is the core of  $H$  in  $G$ , that is, the subgroup of  $H$  generated by all normal subgroups of  $G$  contained in  $H$ .

Let us recall that a subgroup  $H$  of  $G$  is strongly  $\sigma$ -subnormal in  $G$  [6] if  $H^G / H_G$  is  $\sigma$ -nilpotent.

If  $\sigma(H) \subseteq \Pi$ , then  $H$  is called a  $\Pi$ -subgroup of  $G$ . A  $\Pi$ -subgroup  $H$  of  $G$  is called a Hall  $\Pi$ -subgroup of  $G$  if  $\sigma(|G:H|) \cap \Pi = \emptyset$ .

In this paper, we prove the following result.

**Theorem 0.1.** *Let  $G$  be a group. If  $H$  is a Hall  $\Pi$ -subgroup of  $G$ , then the set of all strongly  $\sigma$ -subnormal subgroups of  $G$  which permute with  $H$  forms a sublattice in  $\mathcal{L}(G)$ .*

Taking in Theorem 0.1  $H = G$ , we get from this theorem the following two results.

**Corollary 0.2** (A.N. Skiba [6]). *The set of all strongly  $\sigma$ -subnormal subgroups of  $G$  forms a sublattice in  $\mathcal{L}(G)$ .*

**Corollary 0.3.** *Let  $G$  be a group. If  $H$  is a Hall  $\Pi$ -subgroup of  $G$ , then the set of all strongly  $\sigma$ -subnormal subgroups of  $G$  which permute with  $H$  forms a sublattice in  $\mathcal{L}(G)$ .*

Let us recall that  $G$  is said to be: (i) a  $D_\pi$ -group if  $G$  possesses a Hall  $\pi$ -subgroup  $E$  and every  $\pi$ -subgroup of  $G$  is contained in some conjugate of  $E$ ; (ii) a  $\sigma$ -full group of Sylow type [2] if every subgroup  $E$  of  $G$  is a  $D_{\sigma_i}$ -group for every  $\sigma_i \in \sigma(E)$ .

In view of [8, Theorem 1.2.14], every Sylow permutable subgroup of  $G$  is strongly subnormal in  $G$ . On the other hand, if  $G$  is a  $\sigma$ -full group of Sylow type, then every  $\sigma$ -permutable subgroup is strongly  $\sigma$ -subnormal in  $G$  by Theorem B in [2]. Therefore, since the intersection of any set of sublattices of a lattice is a sublattice of this lattice, we also get from Theorem 0.1 the following two known results.

**Corollary 0.4** (Kegel [9]). *The set of all Sylow permutable subgroups of  $G$  forms a sublattice in  $\mathcal{L}(G)$ .*

**Corollary 0.5** (A.N. Skiba [2]). *If  $G$  is a  $\sigma$ -full group of Sylow type, then the set of all  $\sigma$ -permutable subgroups of  $G$  forms a sublattice in  $\mathcal{L}(G)$ .*

## 1 Lemmas used

**Lemma 1.1** (A.N. Skiba [2]). *The class  $\mathfrak{N}_\sigma$ , of all  $\sigma$ -nilpotent groups, is a hereditary Fitting formation.*

**Lemma 1.2** [5, Ch. A, Proposition 1.6]. *Let  $A, B$  and  $H$  be subgroups of  $G$ . If  $AH = HA$  and  $BH = HB$ , then  $\langle A, B \rangle H = H \langle A, B \rangle$ .*

We use  $O^\Pi(G)$  to denote the subgroup of  $G$  generated by all its  $\Pi'$ -subgroups, where  $\Pi' = \sigma \setminus \Pi$ ;  $O_\Pi(G)$  is the product of all normal  $\Pi$ -subgroups of  $G$ .

**Lemma 1.3** (A.N. Skiba [2, Lemma 2.6]). *If  $A$  is  $\sigma$ -subnormal in  $G$  and  $\sigma(|G:A|) \subseteq \Pi$ -number, then  $O^\Pi(A) = O^\Pi(G)$ .*

## 2 Proof of Theorem 0.1

*Proof.* Let us assume that this theorem is false and let  $G$  be a counterexample of minimal order.

Let  $\mathcal{L}$  be the set of all strongly  $\sigma$ -subnormal subgroups  $L$  of  $G$  which permute with  $H$ .

Let  $A, B \in \mathcal{L}$  and let  $K = \langle A, B \rangle$ ,  $V = A \cap B$ .

First we show that  $K$  is strongly  $\sigma$ -subnormal in  $G$ . By hypothesis,  $A^G / A_G$  is  $\sigma$ -nilpotent. Therefore, in view of the isomorphisms

$$\begin{aligned} A^G(A_G B_G) / A_G B_G &\simeq A^G / (A^G \cap A_G B_G) = \\ &= A^G / A_G(A^G \cap B_G) \simeq \\ &\simeq (A^G / A_G) / (A_G(A^G \cap B_G) / A_G), \end{aligned}$$

we get that

$$A^G(A_G B_G) / A_G B_G \in \mathfrak{N}_\sigma$$

since the class  $\mathfrak{N}_\sigma$  is closed under taking homomorphic images by Lemma 1.1.

Similarly, we can get that

$$B^G(A_G B_G) / A_G B_G \in \mathfrak{N}_\sigma.$$

Moreover,

$$\begin{aligned} A^G B^G / A_G B_G &= \\ &= (A^G(A_G B_G) / A_G B_G)(B^G(A_G B_G) / A_G B_G) \end{aligned}$$

and so, we have

$$A^G B^G / A_G B_G \in \mathfrak{N}_\sigma$$

since the class  $\mathfrak{N}_\sigma$  is a Fitting formation by Lemma 1.1.

Next note that  $\langle A, B \rangle^G = A^G B^G$  and  $A_G B_G \leq \langle A, B \rangle_G$ . Therefore we get that

$$\langle A, B \rangle^G / \langle A, B \rangle_G \in \mathfrak{N}_\sigma$$

since the class  $\mathfrak{N}_\sigma$  is closed under taking homomorphic images by Lemma 1.1. Hence  $\langle A, B \rangle$  is strongly  $\sigma$ -subnormal in  $G$ .

Moreover, in view of Lemma 1.2,

$$\langle A, B \rangle H = H \langle A, B \rangle$$

since  $AH = HA$  and  $BH = HB$  by the choice of  $A$  and  $B$ . Therefore  $K \in \mathcal{L}$ .

Now we show that  $V \in \mathcal{L}$ . First note that

$$(A \cap B)_G = A_G \cap B_G.$$

On the other hand, from the isomorphism

$$\begin{aligned} (A^G \cap B^G) / (A_G \cap B^G) &= \\ &= (A^G \cap B^G) / (A_G \cap B^G \cap A^G) \simeq \\ &\simeq A_G (B^G \cap A^G) / A_G \leq A^G / A_G \end{aligned}$$

we get that

$$(A^G \cap B^G) / (A_G \cap B^G) \in \mathfrak{N}_\sigma$$

since the class  $\mathfrak{N}_\sigma$  is closed under taking normal subgroup by Lemma 1.1. Similarly, we get that

$$(B^G \cap A^G) / (B_G \cap A^G) \in \mathfrak{N}_\sigma.$$

But then we get that

$$(A^G \cap B^G) / (A_G \cap B_G) \in \mathfrak{N}_\sigma$$

since the class  $\mathfrak{N}_\sigma$  is a formation by Lemma 1.1.

It is also clear that

$$(A \cap B)^G \leq A^G \cap B^G.$$

Therefore we get that

$$(A \cap B)^G / (A \cap B)_G \in \mathfrak{N}_\sigma.$$

Therefore  $A \cap B$  is strongly  $\sigma$ -subnormal in  $G$ .

Finally, we show that  $V = A \cap B$  is permutable with  $H$ . Let us assume that this is false. Then  $G$  is not a  $\Pi$ -group, since otherwise we have  $H = G$  and so

$$G = (A \cap B)H = H(A \cap B).$$

First, let us assume that  $R := (A \cap B)_G \neq 1$ . Then

$$(A/R)^G / (A/R)_G = (A^G/R) / (A_G/R) \simeq A^G / A_G$$

is  $\sigma$ -nilpotent, so  $A/R$  is strongly  $\sigma$ -subnormal in  $G/R$ . Similarly,  $B/R$  is strongly  $\sigma$ -subnormal in  $G/R$ . It is also clear that  $HR/R$  is a Hall  $\Pi$ -subgroup of  $G/R$  and  $A/R$  and  $B/R$  permute with  $HR/R$ , so the choice of  $G$  implies that

$$\begin{aligned} ((A \cap B)/R)(HR/R) &= \\ &= ((A/R) \cap (B/R))(HR/R) = \\ &= (HR/R)((A/R) \cap (B/R)) = \\ &= (HR/R)((A \cap B)/R). \end{aligned}$$

But then

$$(A \cap B)H = (A \cap B)HR = HR(A \cap B) = H(A \cap B),$$

which is a contradiction.

Thus,  $(A \cap B)_G = 1$ , so  $(A \cap B)^G$  is  $\sigma$ -nilpotent and hence  $(A \cap B)^G = V \times W$ , where  $W$  is a Hall  $\Pi$ -subgroup of  $(A \cap B)^G$ . Then  $W \leq H$ . It is also clear that  $A \cap B = L \times K$ , where  $K$  is a Hall  $\Pi$ -subgroup of  $A \cap B$  and that  $K \leq H$ . Moreover,

$$L = O^\Pi(A \cap B) = O_{\Pi'}(A \cap B).$$

Now we show that  $H \leq N_G(L)$ . Indeed, we have  $H \leq N_G(O^\Pi(A))$  and  $H \leq N_G(O^\Pi(B))$  by Lemma 1.3, so  $H \leq N_G(O^\Pi(A) \cap O^\Pi(B))$ .

Now observe that  $O^\Pi(A) \cap O^\Pi(B)$  is normal in  $A \cap B$  and from

$$(A \cap B) / (A \cap O^\Pi(A) \cap B) \simeq (A \cap B) O^\Pi(A) / O^\Pi(A)$$

and

$$(A \cap B) / (B \cap O^\Pi(B) \cap A) \simeq (A \cap B) O^\Pi(B) / O^\Pi(B)$$

we get that

$$(A \cap B) / (O^\Pi(A) \cap O^\Pi(B)) =$$

$$= (A \cap B) / ((A \cap O^\Pi(A) \cap B) \cap (B \cap O^\Pi(B) \cap A))$$

is a  $\Pi$ -group. Hence

$$L = O_{\Pi'}(A \cap B) = O_{\Pi'}(O^\Pi(A) \cap O^\Pi(B)),$$

so  $H \leq N_G(L)$ .

Since  $A \cap B = L \times K$ , where  $K \leq H$  and  $H \leq N_G(L)$ , we have

$$\begin{aligned} (A \cap B)H &= (L \times K)H = LH = HL = \\ &= H(L \times K) = H(A \cap B), \end{aligned}$$

a contradiction. Therefore  $V \in \mathcal{L}$ , so  $\mathcal{L}$  is a sublattice of the lattice  $\mathcal{L}(G)$ .  $\square$

## REFERENCES

1. Skiba, A.N. On  $\sigma$ -properties of finite groups I / A.N. Skiba // Problems of Physics, Mathematics and Technics. – 2014. – № 4 (21). – P. 89–96.
2. Skiba, A.N. On  $\sigma$ -subnormal and  $\sigma$ -permutable subgroups of finite groups / A.N. Skiba // J. Algebra. – 2015. – № 436. – P. 1–16.
3. Skiba, A.N. A generalization of a Hall theorem / A.N. Skiba // J. Algebra Appl. – 2015. – Vol. 15, № 4. – P. 21–36.
4. Skiba, A.N. On Some Results in the Theory of Finite Partially Soluble Groups / A.N. Skiba // Commun. Math. Stat. – 2016. – № 4. – P. 281–309.
5. Doerk, K. Finite soluble groups / K. Doerk, T. Hawkes. – Berlin – New York: Walter de Gruyter, 1992.
6. Skiba, A.N. On sublattices of the subgroup lattice defined by formation Fitting sets / A.N. Skiba // J. Algebra. – 2020. – № 550. – P. 69–85.
7. Shemetkov, L.A. Formations of Algebraic Systems / L.A. Shemetkov, A.N. Skiba // Moscow: Nauka, 1989.
8. Ballester-Bolinches, A. Products of Finite Groups / A. Ballester-Bolinches, R. Esteban-Romero, M. Asaad. – Berlin – New York: Walter de Gruyter, 2010.
9. Kegel, O.H. Sylow-Gruppen und Subnormalteilerendlicher Gruppen / O.H. Kegel // Math. Z. – 1962. – № 78. – P. 205–221.

The article was submitted 12.09.2025.

## Информация об авторах

Косенок Николай Сергеевич – к.ф.-м.н., доцент  
Близнец Игорь Васильевич – к.ф.-м.н., доцент