

ЧЕБЫШЕВСКИЙ СБОРНИК

Том 20. Выпуск 2.

УДК 512.542

DOI 10.22405/2226-8383-2019-20-2-391-398

О дисперсивных по Оре \mathfrak{F} -гиперцентральных подгруппах конечных групп

В. И. Мурашко

Мурашко Вячеслав Игоревич — магистрант, Гомельский государственный университет им. Ф. Скорины (г. Гомель, Республика Беларусь).

e-mail: mvimath@yandex.ru

Аннотация

Рассматриваются только конечные группы. Пусть A — группа автоморфизмов группы G , содержащая все внутренние автоморфизмы, и F — максимальный внутренний локальный экран насыщенной формации \mathfrak{F} . A -композиционный фактор H/K группы G называется A - \mathfrak{F} -центральным, если $A/C_A(H/K) \in F(p)$ для всех $p \in \pi(H/K)$. A - \mathfrak{F} -гиперцентром G называется наибольшая A -допустимая подгруппа G , все A -композиционные факторы ниже которой A - \mathfrak{F} -центральны. Обозначается $Z_{\mathfrak{F}}(G, A)$.

Напомним, что группа G называется дисперсивной по Оре, если G имеет нормальную холлову $\{p_1, \dots, p_i\}$ -подгруппу для $1 \leq i \leq n$, где $p_1 > \dots > p_n$ — все простые делители $|G|$. Главным результатом работы является: Пусть \mathfrak{F} — наследственная насыщенная формация, F — её максимальный внутренний локальный экран и N — дисперсивная по Оре A -допустимая подгруппа группы G , где $\text{Inn}G \leq A \leq \text{Aut}G$. Тогда и только тогда $N \leq Z_{\mathfrak{F}}(G, A)$, когда $N_A(P)/C_A(P) \in F(p)$ для любых силовской p -подгруппы P группы N и простого делителя p порядка N .

В качестве следствий были получены известные результаты Р. Бэра о нормальных подгруппах в сверхразрешимом гиперцентре и элементах гиперцентра.

Пусть G — группа. Напомним, что

$$L_n(G) = \{x \in G \mid [x, \alpha_1, \dots, \alpha_n] = 1 \forall \alpha_1, \dots, \alpha_n \in \text{Aut}G\}$$

и G называется автонильпотентной, если $G = L_n(G)$ для некоторого натурального n . Из главного результата можно извлечь критерии автонильпотентности групп. В частности, группа G автонильпотентна тогда и только тогда, когда она является прямым произведением своих силовских подгрупп и группа автоморфизмов любой силовской p -подгруппы группы G является p -группой для любого простого делителя p порядка G . Приведены примеры автонильпотентных групп нечетного порядка.

Ключевые слова: Конечная группа, нильпотентная группа, сверхразрешимая группа, автонильпотентная группа, A - \mathfrak{F} -гиперцентр группы, наследственная насыщенная формация.

Библиография: 25 названий.

Для цитирования:

В. И. Мурашко. О дисперсивных по Оре \mathfrak{F} -гиперцентральных подгруппах конечных групп // Чебышевский сборник, 2019, т. 20, вып. 2, с. 391–398.

CHEBYSHEVSKII SBORNIK

Vol. 20. No. 2.

UDC 512.542

DOI 10.22405/2226-8383-2019-20-2-391-398

On the \mathfrak{F} -hypercentral subgroups with the sylow tower property of finite groups

V. I. Murashka

Murashka Viachaslau Igaravich — graduate student, Francisk Skorina Gomel State University (Gomel, Republic of Belarus).

e-mail: mvimath@yandex.ru

Abstract

Throughout this paper all groups are finite. Let A be a group of automorphisms of a group G that contains all inner automorphisms of G and F be the canonical local definition of a saturated formation \mathfrak{F} . An A -composition factor H/K of G is called A - \mathfrak{F} -central if $A/C_A(H/K) \in F(p)$ for all $p \in \pi(H/K)$. The A - \mathfrak{F} -hypercenter of G is the largest A -admissible subgroup of G such that all its A -composition factors are A - \mathfrak{F} -central. Denoted by $Z_{\mathfrak{F}}(G, A)$.

Recall that a group G satisfies the Sylow tower property if G has a normal Hall $\{p_1, \dots, p_i\}$ -subgroup for all $1 \leq i \leq n$ where $p_1 > \dots > p_n$ are all prime divisors of $|G|$. The main result of this paper is: Let \mathfrak{F} be a hereditary saturated formation, F be its canonical local definition and N be an A -admissible subgroup of a group G where $\text{Inn}G \leq A \leq \text{Aut}G$ that satisfies the Sylow tower property. Then $N \leq Z_{\mathfrak{F}}(G, A)$ if and only if $N_A(P)/C_A(P) \in F(p)$ for all Sylow p -subgroups P of N and every prime divisor p of $|N|$.

As corollaries we obtained well known results of R. Baer about normal subgroups in the supersoluble hypercenter and elements in the hypercenter.

Let G be a group. Recall that $L_n(G) = \{x \in G \mid [x, \alpha_1, \dots, \alpha_n] = 1 \forall \alpha_1, \dots, \alpha_n \in \text{Aut}G\}$ and G is called autonilpotent if $G = L_n(G)$ for some natural n . The criteria of autonilpotency of a group also follow from the main result. In particular, a group G is autonilpotent if and only if it is the direct product of its Sylow subgroups and the automorphism group of a Sylow p -subgroup of G is a p -group for all prime divisors p of $|G|$. Examples of odd order autonilpotent groups were given.

Keywords: Finite group, nilpotent group, supersoluble group, autonilpotent group, A - \mathfrak{F} -hypercenter of a group, hereditary saturated formation.

Bibliography: 25 titles.

For citation:

V. I. Murashka, 2019, "On the \mathfrak{F} -hypercentral subgroups with the sylow tower property of finite groups", *Chebyshevskii sbornik*, vol. 20, no. 2, pp. 391–398.

1. Introduction and results

Throughout this paper all groups are finite and G always denotes a finite group. Recall that $\text{Aut}G$ and $\text{Inn}G$ are the groups of all and inner automorphisms of G respectively.

Let A be a group of automorphisms of G . Kaloujnine [1] and Hall [2] showed that if A stabilizes some chain of subgroups of G , then A is nilpotent. Huppert [3] and Shemetkov [4] showed that if G has A -admissible series with prime indexes, then A is supersoluble. Shemetkov [4] and Schmid [5] obtained analogues results for a solubly saturated formation \mathfrak{F} . Note that the \mathfrak{F} -hypercenter of G with respect to A played an important role in their research.

Recall that $\pi(G)$ is the set of all prime divisors of $|G|$. A *formation* is a class \mathfrak{F} of groups with the following properties: (a) every homomorphic image of an \mathfrak{F} -group is an \mathfrak{F} -group, and (b) if G/M and G/N are \mathfrak{F} -groups, then $G/(M \cap N) \in \mathfrak{F}$. A formation \mathfrak{F} is said to be: *saturated* if $G \in \mathfrak{F}$ whenever $G/\Phi(G) \in \mathfrak{F}$ where $\Phi(G)$ is the Frattini's subgroup of G ; *hereditary* if $H \in \mathfrak{F}$ whenever $H \leq G \in \mathfrak{F}$. A function of the form $f : \mathbb{P} \rightarrow \{\text{formations}\}$ is called a *formation function*. Recall [6, p. 356] that a formation \mathfrak{F} is called *local* if $\mathfrak{F} = (G \mid G/C_G(H/K) \in f(p) \text{ for every } p \in \pi(H/K) \text{ and every chief factor } H/K \text{ of } G)$ for some formation function f . In this case f is called a *local definition* of \mathfrak{F} . By the Gaschütz-Lubeseder-Schmid theorem, a formation is local if and only if it is non-empty and saturated. Recall that if \mathfrak{F} is a local formation, there exists a unique formation function F , defining \mathfrak{F} , such that $F(p) = \mathfrak{N}_p F(p) \subseteq \mathfrak{F}$ for every $p \in \mathbb{P}$ by Proposition 3.8 [6, p. 360]. In this case F is called the *canonical local definition* of \mathfrak{F} .

Let $\text{Inn}G \leq A$ be a group of automorphisms of G and F be the canonical local definition of a local formation \mathfrak{F} . An A -composition factor H/K of G is called *A - \mathfrak{F} -central* if $A/C_A(H/K) \in F(p)$ for all $p \in \pi(H/K)$. The *A - \mathfrak{F} -hypercenter* of G is the largest A -admissible subgroup of G such that all its A -composition factors are A - \mathfrak{F} -central. This subgroup always exists by Lemma 6.4 [6, p. 387]. It is denoted by $Z_{\mathfrak{F}}(G, A)$. If $A = \text{Inn}G$, then it is just the \mathfrak{F} -hypercenter $Z_{\mathfrak{F}}(G)$ of G . If $\mathfrak{F} = \mathfrak{N}$ is the class of all nilpotent groups, then we use $Z_{\infty}(G, A)$ to denote the A -hypercenter $Z_{\mathfrak{N}}(G, A)$ of G . Recently the subgroups of $Z_{\infty}(G, A)$ have been studied for example in [7, 8, 9, 10, 11].

Recall that $\text{Syl}_p G$ is the set of all Sylow subgroups of G ; G *satisfies the Sylow tower property* if G has a normal Hall $\{p_1, \dots, p_i\}$ -subgroup for all $1 \leq i \leq n$ where $p_1 > \dots > p_n$ are all prime divisors of $|G|$. It is well known that a supersoluble group satisfies the Sylow tower property. Recently series of hereditary saturated formations of groups that satisfy the Sylow tower property have been constructed (see, [12, 13, 14, 15]).

THEOREM 1. *Let \mathfrak{F} be a hereditary saturated formation, F be its canonical local definition and N be an A -admissible subgroup of G where $\text{Inn}G \leq A \leq \text{Aut}G$ that satisfies the Sylow tower property. Then $N \leq Z_{\mathfrak{F}}(G, A)$ if and only if $N_A(P)/C_A(P) \in F(p)$ for all $P \in \text{Syl}_p(N)$ and $p \in \pi(N)$.*

Author obtained particular cases of this theorem for $A = \text{Inn}G$ and two formations of supersoluble type in [16, 17]. Recall that a group G is called *strictly p -closed* if $G/O_p(G)$ is abelian of exponent dividing $p - 1$. We use \mathfrak{U} to denote the class of all supersoluble groups.

COROLLARY 1 (R. Baer [18]). *Let N be a normal subgroup of G . Then $N \leq Z_{\mathfrak{U}}(G)$ if and only if N satisfies the Sylow tower property and $N_G(P)/C_G(P)$ is strictly p -closed for all $P \in \text{Syl}_p(N)$ and $p \in \pi(N)$.*

M. R. R. Moghaddam and M. A. Rostamyari (see [9]) introduced the concept of autonilpotent group. Let $L_n(G) = \{x \in G \mid [x, \alpha_1, \dots, \alpha_n] = 1 \forall \alpha_1, \dots, \alpha_n \in \text{Aut}G\}$. Then G is called *autonilpotent* if $G = L_n(G)$ for some natural n . Some properties of autonilpotent groups were studied in [9]. In [8] all abelian autonilpotent groups were described. In particular, abelian autonilpotent non-unit groups of odd order don't exist. It was shown that if a p -group G is autonilpotent, then $\text{Aut}G$ is a p -group (Theorem 2.2 [10]). In [11, p. 45] it was asked: "Does there exist any odd order autonilpotent group?"

COROLLARY 2. *Let p be a prime. A p -group G is autonilpotent if and only if $\text{Aut}G$ is a p -group.*

An example of a p -group G of order p^5 ($p > 3$) such that $\text{Aut}G$ is also a p -group was constructed in [19]. In the library of small groups of GAP [20] there are 30 groups of order 3^6 such that their automorphism groups are also 3-groups (for example groups [729, 31], [729, 41] and [729, 46]). Hence the answer on the question from [11] is positive. From Theorem 2.3 [9] and Lemma 2.9 [10] it follows that a group is autonilpotent iff it is the direct product of its autonilpotent Sylow subgroups.

COROLLARY 3. *A group G is autonilpotent if and only if it is the direct product of its Sylow subgroups and the automorphism group of a Sylow p -subgroup of G is a p -group for all $p \in \pi(G)$.*

The proof of Corollary 3 doesn't use results from [9, 10]. The following result gives the description of elements in $Z_\infty(G, A)$.

COROLLARY 4. *Let g be a p -element of a group G and $\text{Inn}G \leq A \leq \text{Aut}G$. Then $g \in Z_\infty(G, A)$ if and only if $g^\alpha = g$ for every p' -element α of A .*

COROLLARY 5 (R. Baer [21]). *Let p be a prime and G be a group. Then a p -element g of G belongs to $Z_\infty(G)$ if and only if it permutes with all p' -elements of G .*

COROLLARY 6. *A group G is autonilpotent if and only if every automorphism α of G fixes all elements of G whose orders are coprime to the order of α .*

According to Frobenius p -nilpotency criterion (see Theorem 5.26 [22, p. 171]) a group G is nilpotent if and only if $N_G(P)/C_G(P)$ is a p -group for every p -subgroup P of G and every $p \in \pi(G)$.

COROLLARY 7. *A group G is autonilpotent if and only if $N_{\text{Aut}G}(P)/C_{\text{Aut}G}(P)$ is a p -group for every p -subgroup P of G and every $p \in \pi(G)$.*

2. Proves of the results

LEMMA 1 (Lemma 3.6 [16]). *Let P be a p -subgroup of a group G and R be a normal r -subgroup of G where $r \neq p$ are primes. Then $N_G(P)R/R = N_{G/R}(PR/R)$, $C_G(P)R/R = C_{G/R}(PR/R)$ and $N_G(P)/C_G(P) \simeq N_{G/R}(PR/R)/C_{G/R}(PR/R)$.*

PROOF. [Proof of Theorem 1] Let prove Theorem 1 for $A = \text{Inn}G$. In this case $Z_{\mathfrak{F}}(G, A) = Z_{\mathfrak{F}}(G)$.

Sufficiency. Let N be a normal subgroup of $Z_{\mathfrak{F}}(G)$ with the Sylow tower property. So N has a normal Sylow q -subgroup Q . Note that $Q \trianglelefteq G$ and $Q \leq Z_{\mathfrak{F}}(G)$.

Hence $G/C_G(Q) = N_G(Q)/C_G(Q) \in F(q)$ by Lemma 2.5 from [23]. If $Q = N$, then sufficiency is proven.

Let $Q < N$. From $N \leq Z_{\mathfrak{F}}(G)$ it follows that $N/Q \leq Z_{\mathfrak{F}}(G/Q)$. Using induction on the order of G , we may assume that $N_{G/Q}(PQ/Q)/C_{G/Q}(PQ/Q) \in F(p)$ for every $P \in \text{Syl}_p N$ and $p \in \pi(N) \setminus \{q\}$. Hence $N_G(P)/C_G(P) \simeq N_{G/Q}(PQ/Q)/C_{G/Q}(PQ/Q) \in F(p)$ by Lemma 1. Thus sufficiency is proved.

Necessity. Let a group G be a minimal order counterexample with a normal subgroup $N \not\leq Z_{\mathfrak{F}}(G)$ that satisfies the statement of Theorem 1 and p be the greatest prime divisor of $|N|$. Then a Sylow p -subgroup P of N is normal in G . Let H/K be a chief factor of G and $H \leq P$. Since $C_G(P) \leq C_G(H/K)$ and $G/C_G(P) = N_G(P)/C_G(P) \in F(p)$, $G/C_G(H/K) \in F(p)$. It means that $P \leq Z_{\mathfrak{F}}(G)$. Note that $N_{G/P}(RP/P)/C_{G/P}(RP/P) \simeq N_G(R)/C_G(R) \in F(p)$ for every $R \in \text{Syl}_r N$ and $r \in \pi(N) \setminus \{p\}$ by Lemma 1. From $|G/P| < |G|$ it follows that $N/P \leq Z_{\mathfrak{F}}(G/P)$. Thus $N \leq Z_{\mathfrak{F}}(G)$, the contradiction.

Assume now that $\text{Inn}G \leq A \leq \text{Aut}G$. Let $\Gamma = G \rtimes A$. From $\text{Inn}G \leq A$ it follows that groups of automorphisms that are induced by A and Γ on a given section of G are isomorphic. It means that $N_A(H/K)/C_A(H/K) \simeq N_\Gamma(H/K)/C_\Gamma(H/K)$ for a given section H/K of G . In particular, every A -composition A - \mathfrak{F} -central factor of G is an \mathfrak{F} -central chief factor of Γ .

It means that A -admissible subgroup N of G that satisfies the Sylow tower property lies in $Z_{\mathfrak{F}}(G, A)$ if and only if $N \leq Z_{\mathfrak{F}}(\Gamma)$. The later is equivalent to $N_\Gamma(P)/C_\Gamma(P) \simeq N_A(P)/C_A(P) \in F(p)$ for every $P \in \text{Syl}_p N$ and $p \in \pi(N)$. \square

PROOF. [Proof of Corollary 1] Recall that \mathfrak{U} has the canonical local definition F where $F(p)$ is the class of all strictly p -closed groups, every subgroup of $Z_{\mathfrak{U}}(G)$ is supersoluble and every supersoluble group satisfies the Sylow tower property. Thus Corollary 1 directly follows from Theorem 1. \square

PROPOSITION 3. A group G is autonilpotent if and only if $G = Z_\infty(G, \text{Aut}G)$.

PROOF. [Proof] Recall that the canonical local definition of \mathfrak{N} is $F(p) = \mathfrak{N}_p$ where \mathfrak{N}_p is the class of all p -groups. Note that $\text{Aut}G/C_{\text{Aut}G}(L_i(G)/L_{i-1}(G)) \simeq 1 \in F(p)$ for all prime p . Hence every $\text{Aut}G$ -composition factor of G between $L_{i-1}(G)$ and $L_i(G)$ is $\text{Aut}G$ - \mathfrak{N} -central. It means that $L_i(G) \leq Z_\infty(G, \text{Aut}G)$ for every i . So if G is autonilpotent group, then $G = Z_\infty(G, \text{Aut}G)$.

Assume that $G = Z_\infty(G, \text{Aut}G)$. Hence there exists an $\text{Aut}G$ -composition series

$$1 = G_0 \leq \dots \leq G_n = G$$

with $\text{Aut}G$ - \mathfrak{N} -central chief factors. Let $\Gamma = G \rtimes \text{Aut}G$. Note that

$$\text{Aut}G/C_{\text{Aut}G}(G_i/G_{i-1}) \simeq \Gamma/C_\Gamma(G_i/G_{i-1}) \in \mathfrak{N}_p$$

for every $p \in \pi(G_i/G_{i-1})$ by analogy with the proof of Theorem 1. Note that $\Gamma/C_\Gamma(G_i/G_{i-1})$ does not have non-trivial p -subgroups for all $p \in \pi(G_i/G_{i-1})$ by Lemma 3.9 [24, p. 26]. So $\text{Aut}G = C_{\text{Aut}G}(G_i/G_{i-1})$. Hence $[G_i, \text{Aut}G] \leq G_{i-1}$. It means that $[x, \alpha_1, \dots, \alpha_n] = 1$ for all $x \in G$ and $\alpha_1, \dots, \alpha_n \in \text{Aut}G$. Thus $G = L_n(G)$ and G is autonilpotent. \square

PROOF. [Proof of Corollary 3] Note that every nilpotent group satisfies the Sylow tower property and every autonilpotent group is nilpotent.

So a nilpotent group is autonilpotent if and only if $N_{\text{Aut}G}(P)/C_{\text{Aut}G}(P) \in \mathfrak{N}_p$ for every $P \in \text{Syl}_p G$ and $p \in \pi(G)$ by Proposition 3 and Theorem 1. The automorphism group of a direct product of groups was described in [25]. In particular, if $G = P \times H$, where P is a Sylow subgroup of G , then $\text{Aut}G = \text{Aut}P \times \text{Aut}H$ and $N_{\text{Aut}G}(P)/C_{\text{Aut}G}(P) \simeq \text{Aut}P$. \square

PROOF. [Proof of Corollary 2] Directly follows from Corollary 3. \square

PROOF. [Proof of Corollary 4] From $\text{Inn}G \leq A$ it follows that $Z_\infty(G, A) \leq Z_\infty(G)$ is nilpotent. Hence every Sylow subgroup of $Z_\infty(G, A)$ is A -admissible. So

$$N_A(P)/C_A(P) = A/C_A(P) \in \mathfrak{N}_p$$

for every $P \in \text{Syl}_p Z_\infty(G, A)$ and $p \in \pi(Z_\infty(G, A))$ by Theorem 1. It means that if g is a p -element of $Z_\infty(G, A)$, then $g^\alpha = g$ for every p' -element α of A .

Let G_p be the set of all elements of G such that $g^\alpha = g$ for every p' -element α of A and every $g \in G_p$. Note that if $x, y \in G_p$, then $xy \in G_p$. Hence G_p is a subgroup of G . Let $g \in G_p$, $\alpha, \beta \in A$ and β be a p' -element. Then $\beta^{\alpha^{-1}}$ is a p' -element too. Hence $(g^\alpha)^\beta = g^{\alpha\beta\alpha^{-1}} = g^{\beta^{\alpha^{-1}}\alpha} = g^\alpha$. It means that $g^\alpha \in G_p$. Thus G_p is an A -admissible subgroup of G . Let x be a p' -element of G_p . From $\text{Inn}G \leq A$ it follows that $\alpha_x : g \rightarrow g^x$ is a p' -element of A . Hence it acts trivially on G_p . It means that $x \leq Z(G_p)$. Let $P \in \text{Syl}_p G_p$. Then $P \trianglelefteq G_p$. So all p -elements of G_p form a subgroup P . Note that P is A -admissible and $N_A(P)/C_A(P) \in \mathfrak{N}_p$. Therefore $P \leq Z_\infty(G, A)$ by Theorem 1. \square

PROOF. [Proof of Corollary 5] Let g be a p -element of G . Note that $xg = gx$ is equivalent to $g^x = g$ and $\{\alpha_x : g \rightarrow g^x \mid x \text{ is a } p'\text{-element of } G\}$ is the set of all p' -elements of $\text{Inn}G$. Now Corollary 5 directly follows from Corollary 4. \square

PROOF. [Proof of Corollary 6] Directly follows from Proposition 3 and Corollary 4. \square

PROOF. [Proof of Corollary 7] Assume that G is autonilpotent. Then

$$N_{\text{Aut}G}(P)/C_{\text{Aut}G}(P) \in \mathfrak{N}_p$$

for every p -subgroup P of G and $p \in \pi(G)$ by Corollary 6.

Assume now that $N_{\text{Aut}G}(P)/C_{\text{Aut}G}(P) \in \mathfrak{N}_p$ for every p -subgroup P of G and $p \in \pi(G)$. Suppose that G is non-nilpotent. So there is a Schmidt subgroup S of G . Then S has a normal q -subgroup Q for some prime q and there is a q' -element x of S with $x \notin C_S(Q)$ (see Theorem 26.1 [24, p. 243]). Since $\alpha_x : g \rightarrow g^x$ is a non-identity inner automorphism of G of q' -order, $N_{\text{Aut}G}(Q)/C_{\text{Aut}G}(Q) \notin \mathfrak{N}_q$, a contradiction.

Thus G is nilpotent. Hence G is autonilpotent by Proposition 3 and Theorem 1. \square

Acknowledgments

I am grateful to A. F. Vasil'ev for helpful discussions.

REFERENCES

1. Kaloujnine L. 1953, "Über gewisse Beziehungen zwischen einer Gruppe und ihren Automorphismen", Berliner Mathematische Tagung, Berlin, pp. 164–172.
2. Hall P. 1958, "Some sufficient conditions for a group to be nilpotent", *Illinois J. Math.*, vol. 2, no. 4, pp. 787–801.
3. Huppert B. 1954, "Normalteiler und maximale Untergruppen endlicher Gruppen", *Math. Z.*, vol. 60, no. 4, pp. 409–434.
4. Semetkov L. A. 1974, "Graduated formations of groups", *Mathematics of the USSR-Sbornik*, vol. 23, no. 4, pp. 593–611.
5. Schmid P. 1974, "Lokale Formationen endlicher Gruppen", *Math. Z.*, vol. 137, no. 1, pp. 31–48.
6. Doerk K., Hawkes T. 1992, *Finite soluble groups*, Walter de Gruyter, Berlin–New York, 891 p.
7. Hegarty P. V. 1994, "The absolute center of a group", *J. Algebra*, vol. 169, pp. 929–935.
8. Nasrabadi M. M., Gholamiam A. 2014, "On A-nilpotent abelian groups", *Proc. Indian Acad. Sci. (Math. Sci.)*, vol. 124, no. 4, pp. 517–525.
9. Davoudirad S., Moghaddam M. R. R., Rostamyari M. A. 2016, "Autonilpotent groups and their properties", *Asian-European J. Math.*, vol. 9, no. 2, 1650056 (7 pages).
10. Hoseini S., Moghaddam M. R. R., Tajnia S. 2015, "On Auto-nilpotent groups", *Southeast Asian Bull. Math.*, vol. 39, pp. 219–224.
11. Arora H., Karan R. 2018, "On Autonilpotent and Autosoluble Groups", *Note Mat.*, vol. 38, no. 1, pp. 35–45.
12. Vasil'ev A. F., Vasil'eva T. I., Tyutyaynov V. N. 2010, "On the finite groups of supersoluble type", *Sib. Math. J.*, vol. 51, no. 6, pp. 1004–1012.
13. Monakhov V. S., Kniahina V. N. 2013, "Finite groups with \mathbb{P} -subnormal subgroups", *Ricerche mat.*, vol. 62, pp. 307–322.
14. Zimmermann I. 1989, "Submodular subgroups in finite groups", *Math. Z.*, vol. 202, pp. 545–557.
15. Vasilyev V. A. 2015, "Finite groups with submodular Sylow subgroups", *Siberian Math. J.*, vol. 56, no. 6, pp. 1019–1027.
16. Murashka V. I. 2018, "On analogues of Baer's theorems for widely supersoluble hypercenter of finite groups", *Asian-European J. Math.*, vol. 11, no. 3, 1850043 (8 pages).
17. Murashka V. I. 2014, "Properties of the class of finite groups with P-subnormal cyclic primary subgroups", *Dokl. NAN Belarusi*, vol. 58, no. 1, pp. 5–8. (In Russian)
18. Baer R. 1959, "Supersoluble immersion", *Canad. J. Math.*, vol. 11, pp. 353–369.
19. Curran M. J. 1988, "Automorphisms of certain p -groups (p odd)", *Bull. Austral. Math. Soc.*, vol. 38, pp. 299–305.

20. Groups, Algorithms, and Programming (GAP), Version 4.9.1 (2018). Available at <http://www.gap-system.org> (accessed 10 June 2018).
21. Baer R. 1953, "Group Elements of Prime Power Index", Trans. Amer. Math Soc., vol. 75, no. 1, pp. 20–47.
22. Isaacs I. M. 2008, Finite group theory, Graduate studies in mathematics, vol. 92, Providence, American Mathematical Society, 350 p.
23. Skiba A. N. 2012, "On the \mathfrak{F} -hypercentral and the intersection of all \mathfrak{F} -maximal subgroups of a finite group", J. Pure Appl. Algebra, vol. 216, no. 4, pp. 789–799.
24. Shemetkov L. A. 1978, Formations of finite groups, Nauka, Moscow, 272 p. (In Russian)
25. Bidwell J. N. S., Curran M. J., McCaughan D. J. 2006, "Automorphisms of direct products of finite groups", Arch. Math., vol. 86, pp. 481–489.

СПИСОК ЦИТИРОВАННОЙ ЛИТЕРАТУРЫ

1. Kaloujnine L. Über gewisse Beziehungen zwischen einer Gruppe und ihren Automorphismen // Berliner Mathematische Tagung, Berlin, 1953, P. 164–172.
2. Hall P. Some sufficient conditions for a group to be nilpotent // Illinois J. Math. 1958. V. 2, № 4. P. 787–801.
3. Huppert B. Normalteiler und maximale Untergruppen endlicher Gruppen // Math. Z. 1954. V. 60, № 4. P. 409–434.
4. Шеметков Л. А. Ступенчатые формации групп // Матем. сб. 1974. Т. 94(136), № 4(8), С. 628–648.
5. Schmid P. Lokale Formationen endlicher Gruppen // Math. Z. 1974. V. 137, № 1. P. 31–48.
6. Doerk K., Hawkes T. Finite soluble groups. Berlin–New York: Walter de Gruyter, 1992. 891 p.
7. Hegarty P. V. The absolute center of a group // J. Algebra. 1994. V. 169. P. 929–935.
8. Nasrabadi M. M., Gholamiam A. On A-nilpotent abelian groups // Proc. Indian Acad. Sci. (Math. Sci.). 2014. V. 124, № 4. P. 517–525.
9. Davoudirad S., Moghaddam M. R. R., Rostamyari M. A. Autonilpotent groups and their properties // Asian-European J. Math. 2016. V. 9, № 2. 1650056 (7 pages).
10. Hoseini S., Moghaddam M. R. R., Tajnia S. On Auto-nilpotent groups // Southeast Asian Bull. Math. 2015. V. 39. P. 219–224.
11. Arora H., Karan R. On Autonilpotent and Autosoluble Groups // Note Mat. 2018. V. 38, № 1. P. 35–45.
12. Васильев А. Ф., Васильева Т. И., Тютянов В. Н. О конечных группах сверхразрешимого типа // Сиб. мат. журн. 2010. Т. 51, № 6. С. 1270–1281.
13. Monakhov V. S., Kniahina V. N. Finite groups with \mathbb{P} -subnormal subgroups // Ricerche mat. 2013. V. 62, P. 307–322.
14. Zimmermann I. Submodular subgroups in finite groups // Math. Z. 1989. V. 202. P. 545–557.

15. Васильев В. А. Конечные группы с субмодулярными силовскими подгруппами // Сиб. мат. журн. 2015. Т. 56, № 6. С. 1277–1288.
16. Murashka V. I. On analogues of Baer's theorems for widely supersoluble hypercenter of finite groups // Asian-European J. Math. 2018. V. 11, № 3. 1850043 (8 pages).
17. Мурашко, В. И. Свойства класса конечных групп с P -субнормальными циклическими примарными подгруппами // Докл. НАН Беларуси. 2014. Т. 58, № 1. С. 5–8.
18. Baer R. Supersoluble immersion // Canad. J. Math. 1959. V. 11. P. 353–369.
19. Curran M. J. Automorphisms of certain p -groups (p odd) // Bull. Austral. Math. Soc. 1988. V. 38. P. 299–305.
20. Groups, Algorithms, and Programming (GAP), Version 4.9.1. [Электронный ресурс] // URL: <http://www.gap-system.org> (дата обращения 10.06.2018).
21. Baer R. Group Elements of Prime Power Index // Trans. Amer. Math Soc. 1953. V. 75, № 1. P. 20–47.
22. Isaacs I. M. Finite group theory. Graduate studies in mathematics, V. 92. Providence: American Mathematical Society, 2008. 350 p.
23. Skiba A. N. On the \mathfrak{F} -hypercenter and the intersection of all \mathfrak{F} -maximal subgroups of a finite group // J. Pure Appl. Algebra. 2012. V. 216, № 4. P. 789–799.
24. Шеметков Л. А. Формации конечных групп. М.: Наука, 1978. 272 с.
25. Bidwell J. N. S., Curran M. J., McCaughan D. J. Automorphisms of direct products of finite groups // Arch. Math. 2006. V. 86. P. 481–489.

Получено 15.06.2018 г.

Принято в печать 12.07.2019 г.